

Could chronic wounds not heal due to too low local copper levels?

Published 2021

medical hypotheses

This study explores possibility that low levels of copper contribute to the incapacity of the wounds to heal in individuals with diabetic ulcers, decubitus, peripheral vascular, or other wounds with compromised circulation to the wound site.

Chronic wounds are characterized by extensive loss of the integument, clear necrosis, or signs of circulation impairment either localized or more extensive, usually in the limbs, leading to extensive loss of substance.

Contrary to the high susceptibility of microorganisms to copper, human skin is not sensitive to copper and the risk of adverse reactions due to dermal exposure to copper is extremely low.

Several mechanisms for the potent biocidal activity of copper have been proposed, which include alteration of proteins and inhibition of their biological assembly and activity; plasma membrane permeabilization; and membrane lipid peroxidation.

Some observations about the role of copper include:

- Immunohistochemical studies show that wound edges of copper-treated wounds have more prominent VEGF expression
- Copper modulates integrins expressed by suprabasally differentiated keratinocytes during the final healing phase
- GHK, a tripeptide with high affinity for copper ions that was isolated from human plasma, potently reduces tissue oxidative damage after injury and activates tissue remodeling
- Expression of the metallothionein gene is upregulated in the skin following topical application of copper, and in wound margins, particularly in regions of high mitotic activity.

“...in ischemic patients, with poor local copper supply by the circulatory system, exterior supplement of the copper by elution from copper containing dressings, directly to the wounds, will result in healing where conventional treatments fail.”

Gadi Borkow, Jeffrey Gabbay, Richard C. Zatcoff, Could chronic wounds not heal due to too low local copper levels?, Medical Hypotheses, Volume 70, Issue 3, 2008, 70(3): 610-613.

Read the full text online:

<https://www.sciencedirect.com/science/article/abs/pii/S0306987707004124>



ELSEVIER

medical
hypotheses<http://intl.elsevierhealth.com/journals/mehy>

Could chronic wounds not heal due to too low local copper levels?

Gadi Borkow ^{a,*}, Jeffrey Gabbay ^a, Richard C. Zatcoff ^b

^a Cupron Inc., P.O. Box 10973, Greensboro, NC 27404, USA

^b Upstate Podiatry, 727 SE Main Street, Ste. 380, Simpsonville, SC 29681, USA

Received 5 June 2007; accepted 7 June 2007

Summary Copper is an essential trace element involved in numerous human physiological and metabolic processes. It plays a key role in many of the processes that together comprise wound healing, including induction of endothelial growth factor, angiogenesis and expression and stabilization of extracellular skin proteins. We hypothesize that in individuals with diabetic ulcers, decubitus, peripheral vascular, or other wounds which might have compromised circulation to the wound site, that part of the incapacity of the wounds to heal is due to low local copper levels. Contamination of wounds is also an important factor causing impaired wound healing. Importantly, copper has potent broad biocidal properties. In contrast, the risk of adverse skin reactions due to exposure to copper is extremely low. We thus hypothesize that introducing copper into wound dressings would not only reduce the risk of wound and dressing contamination, as silver does but, more importantly, would stimulate faster wound repair directly. This would be done by the release of copper from the wound dressings directly into the wound site inducing angiogenesis and skin regeneration.

© 2007 Elsevier Ltd. All rights reserved.

Introduction

Wounds normally heal in a very orderly, efficient and highly controlled method of repair distinguished by four distinct, but overlapping, phases: hemostasis, inflammation, proliferation and remodelling [1,2]. These highly complex phases, characterized by high metabolic activity at the wound site, demand a fluent supply of oxygen

and nutrients. The altered microenvironment found in wounds, such as low pH, reduced oxygen tension and increased lactate, actually initiate the release of factors by epidermal cells, fibroblasts, macrophages and vascular endothelial cells needed to bring in new blood supply. This process of neovascularisation is stimulated by vascular endothelial cell growth factor (VEGF), basic fibroblast growth factor (bFGF) and transforming growth factor (TGF)- β . Fibroblasts are responsible for producing the new matrix needed to restore structure and function to the injured tissue. Fibroblasts attach to fibrin and integrin cables and begin to produce collagen, which then becomes cross-linked [3,4].

* Corresponding author. Address: Hameyasdim 44, Kfar Gibton, 76910, Israel. Tel.: +972 546 611287/+1 866 317 4670; fax: +972 8 9491254.

E-mail address: gadi@cupron.com (G. Borkow).

Ischemia in chronic wounds

The events of early wound healing reflect a finely balanced environment leading to uncomplicated and rapid wound healing. Chronic wounds or "complex wounds" [5] which, for many reasons, have lost this fine balance, are usually associated with systemic pathologies that impair normal healing causing wounds to fail to heal with simple care and requiring special attention. Chronic wounds are characterized by extensive loss of the integument, clear necrosis, or signs of circulation impairment either localized or more extensive, usually in the limbs, leading to extensive loss of substance. A key characteristic common to chronic wounds is ischemia, due to microangiopathic disease in diabetic patients [6], chronic venous insufficiency in chronic venous ulcers [7], vasculitis in patients with autoimmune disease or under immunosuppressive drug therapy [8] or necrosis of the integument in pressure sores [9].

Involvement of systemic copper in wound healing

The function of many factors involved in the efficient and highly controlled repair wound healing mechanism is dependant on their interaction with copper. The following supports this notion:

- (i) Inducible VEGF expression is sensitive to copper [10]. Among many known growth factors, VEGF is believed to be the most prevalent, efficacious, and long-term signal that is known to stimulate angiogenesis in wounds. In both *in vitro* and *in vivo* models, CuSO_4 clearly promotes angiogenic responses. Topical copper sulphate accelerated the closure of excisional murine dermal wounds. Histological analysis of wound-edge tissue substantiated that CuSO_4 treatment not only accelerated wound closure but that the quality of regenerating tissue was distinctly different. CuSO_4 treatment was associated with more hyperproliferative epithelial tissue and the density of cells in the granulation layer of copper-treated wounds was clearly higher. Immunohistochemical studies show that wound edges of copper-treated wounds have more prominent VEGF expression [10].
- (ii) The export of fibroblast growth factor (FGF)1 and interleukin (IL)-1 α , two pro-angiogenic polypeptides, is based on the Cu^{2+} -dependent formation of multiprotein complexes containing the S100A13 protein.

The formation of a such a multiprotein aggregate enables the release of FGF1 in response to stress [11].

- (iii) Furthermore, copper chelation represses the vascular response to injury [12]. These observations have also led to the development of anti-copper-based, anti-angiogenic strategies for the treatment of cancer [13,14].
- (iv) Integrins expressed by basal layer keratinocytes play an essential part in wound healing. Copper modulates integrins expressed by suprabasally differentiated keratinocytes during the final healing phase [15].
- (v) The stability of fibronectin mats (Fn-mat), which have potential applications in tissue engineering, is increased by treating them with micromolar concentrations of copper ions [16]. Low concentrations of copper (1 μM) not only caused significant fibronectin stabilisation but the greatest amount of cell ingrowth was observed for copper treated cables [17].
- (vi) Expression of the metallothionein gene is up-regulated in the skin following topical application of copper, and in wound margins, particularly in regions of high mitotic activity. The action of metallothioneins in these processes may result from the large number of zinc-dependent and copper-dependent enzymes required for cell proliferation and matrix remodelling [18].
- (vii) GHK, a tripeptide with high affinity for copper ions that was isolated from human plasma, potentially reduces tissue oxidative damage after injury and activates tissue remodelling [19]. Topical application of GHK resulted in accelerated wound healing in ischemic open wounds [20]. Interestingly, the increase in metalloproteinase-2 levels in conditioned media of cultured fibroblasts GHK-Cu(2+), could be reproduced by copper ions alone but not by the tripeptide GHK alone [21], indicating the crucial role of copper in the remodelling of extracellular matrix, a central step in wound repair. Similarly, the copper tripeptide accelerates the growth of normal and irradiated fibroblasts to the point where treated irradiated fibroblasts approximate the population-doubling time of normal controls [22]. Accordingly, it was found that GHK incorporated collagen increases the copper concentration by nine fold at the wound site indicating the wound healing property of GHK can also be linked with copper localization [23].

As delineated above, copper ions play an important role especially, but not only, during

the proliferation and remodelling phases. It is therefore not surprising that many over-the-counter treatments for wound healing contain copper [24,25]. The importance of copper in wound healing is further demonstrated by the positive effect of its administration in cases of severe burn trauma in children [26] and in the management of phosphorus burns [27].

Copper and wound infections

It is established that infections may delay healing, cause failure of healing, and even cause wound deterioration [28]. Microbial pathogens delay wound healing through several different mechanisms: persistent production of inflammatory mediators; metabolic wastes; toxins; tissue hypoxia; rendering the granulation tissue hemorrhagic and fragile; reducing fibroblast number and collagen production; damaging reepithelization [29,30]; maintaining the activated state of neutrophils, which produce cytolytic enzymes and free oxygen radicals, and competing with host cells for nutrients and oxygen necessary for wound healing [31]. There is increasing evidence that bacteria within chronic wounds live within biofilm communities, in which the bacteria are protected from host defences and develop resistance to antibiotic treatment [28]. Thus, keeping the wound aseptic or maintaining a low bioburden is an important aim in wound healing, especially when dealing with chronic wounds.

Copper ions, either alone or in copper complexes, have been used for centuries to disinfect liquids, solids and human tissue. Several mechanisms for the potent biocidal activity of copper have been proposed, which include alteration of proteins and inhibition of their biological assembly and activity; plasma membrane permeabilization; and membrane lipid peroxidation [32]. In contrast to the resistant microbes that have evolved to antibiotics in less than 50 years of use, tolerant microbes to copper are extremely rare even though copper has been a part of the earth for millions of years. This lack of resistance to copper may be explained by the capacity of copper to damage in parallel many key factors in micro-organisms [32].

Contrary to the high susceptibility of micro-organisms to copper, human skin is not sensitive to copper and the risk of adverse reactions due to dermal exposure to copper is extremely low [33,34]. Copper is not only considered safe to humans, as demonstrated by the widespread and prolonged use by women of copper intrauterine devices (IUDs) [35,36], but it is an essential metal

needed for normal metabolic processes. The National Academy of Sciences Committee established the US recommended Daily Allowance of 0.9 mg of copper for normal adults [37].

The hypothesis

We hypothesize that in diabetic ulcers, decubitus, and other wounds that occur in individuals with potentially compromised blood supply to the wound site, part of the incapacity of the wounds to heal is due to low local copper levels, which occur due to the compromised blood circulation.

Taking together the potent biocidal activities of copper and its roles in the wound healing process, strongly supports the notion that the addition or application of copper or copper containing products, such as Band-Aids and gauze containing copper, to wounds, may significantly enhance the wound healing process. Furthermore, in ischemic patients, with poor local copper supply by the circulatory system, exterior supplement of the copper by elution from copper containing dressings, directly to the wounds, will result in healing where conventional treatments fail.

References

- [1] Richardson M. Acute wounds: an overview of the physiological healing process. *Nurs Times* 2004;100:50–3.
- [2] Diegelmann RF, Evans MC. Wound healing: an overview of acute, fibrotic and delayed healing. *Front Biosci* 2004;9: 283–9.
- [3] Sorrell JM, Baber MA, Caplan AI. Clonal characterization of fibroblasts in the superficial layer of the adult human dermis. *Cell Tissue Res* 2007;327:499–510.
- [4] Gailit J, Clarke C, Newman D, Tonnesen MG, Mosesson MW, Clark RA. Human fibroblasts bind directly to fibrinogen at RGD sites through integrin $\alpha(v)\beta 3$. *Exp Cell Res* 1997;232:118–26.
- [5] Ferreira MC, Tuma Jr P, Carvalho VF, Kamamoto F. Complex wounds. *Clinics* 2006;61:571–8.
- [6] Bowering CK. Diabetic foot ulcers. Pathophysiology, assessment, and therapy. *Can Fam Physician* 2001;47:1007–16.
- [7] Abbade LP, Lastoria S. Venous ulcer: epidemiology, pathophysiology, diagnosis and treatment. *Int J Dermatol* 2005;44:449–56.
- [8] Coelho S, Amarello M, Ryan S, Reddy M, Sibbald RG. Rheumatoid arthritis-associated inflammatory leg ulcers: a new treatment for recalcitrant wounds. *Int Wound J* 2004;1:81–4.
- [9] Cullum N, Nelson EA, Nixon J. Pressure ulcers. *Clin Evid* 2006;2592–606.
- [10] Sen CK, Khanna S, Venojarvi M, et al. Copper-induced vascular endothelial growth factor expression and wound healing. *Am J Physiol Heart Circ Physiol* 2002;282: H1821–7.
- [11] Prudovsky I, Mandinova A, Soldi R, et al. The non-classical export routes: FGF1 and IL-1 α point the way. *J Cell Sci* 2003;116:4871–81.

- [12] Mandinov L, Mandinova A, Kyurkchiev S, et al. Copper chelation represses the vascular response to injury. *Proc Natl Acad Sci USA* 2003;100:6700–5.
- [13] Mandinova A, Soldi R, Graziani I, et al. S100A13 mediates the copper-dependent stress-induced release of IL-1 α from both human U937 and murine NIH 3T3 cells. *J Cell Sci* 2003;116:2687–96.
- [14] Voelker R. Copper and cancer. *JAMA* 2000;283:994.
- [15] Tenaud I, Sainte-Marie I, Jumbou O, Litoux P, Dreno B. In vitro modulation of keratinocyte wound healing integrins by zinc, copper and manganese. *Br J Dermatol* 1999;140: 26–34.
- [16] Ahmed Z, Idowu BD, Brown RA. Stabilization of fibronectin mats with micromolar concentrations of copper. *Biomaterials* 1999;20:201–9.
- [17] Ahmed Z, Briden A, Hall S, Brown RA. Stabilisation of cables of fibronectin with micromolar concentrations of copper: in vitro cell substrate properties. *Biomaterials* 2004;25: 803–12.
- [18] Lansdown AB. Metallothioneins: potential therapeutic aids for wound healing in the skin. *Wound Repair Regen* 2002;10:130–2.
- [19] Maquart FX, Bellon G, Chaqour B, et al. In vivo stimulation of connective tissue accumulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺ in rat experimental wounds. *J Clin Invest* 1993;92:2368–76.
- [20] Canapp Jr SO, Farese JP, Schultz GS, et al. The effect of topical tripeptide-copper complex on healing of ischemic open wounds. *Vet Surg* 2003;32:515–23.
- [21] Simeon A, Emonard H, Hornebeck W, Maquart FX. The tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺ stimulates matrix metalloproteinase-2 expression by fibroblast cultures. *Life Sci* 2000;67:2257–65.
- [22] Pollard JD, Quan S, Kang T, Koch RJ. Effects of copper tripeptide on the growth and expression of growth factors by normal and irradiated fibroblasts. *Arch Facial Plast Surg* 2005;7:27–31.
- [23] Arul V, Gopinath D, Gomathi K, Jayakumar R. Biotinylated GHK peptide incorporated collagenous matrix: a novel biomaterial for dermal wound healing in rats. *J Biomed Mater Res B Appl Biomater* 2005;73:383–91.
- [24] Pereira CE, Felcman J. Correlation between five minerals and the healing effect of Brazilian medicinal plants. *Biol Trace Elem Res* 1998;65:251–9.
- [25] Schlemm DJ, Crowe MJ, McNeill RB, Stanley AE, Keller SJ. Medicinal yeast extracts. *Cell Stress Chaperones* 1999;4: 171–6.
- [26] Cunningham JJ, Leffell M, Harmatz P. Burn severity, copper dose, and plasma ceruloplasmin in burned children during total parenteral nutrition. *Nutrition* 1993;9:329–32.
- [27] Chou TD, Lee TW, Chen SL, et al. The management of white phosphorus burns. *Burns* 2001;27:492–7.
- [28] Edwards R, Harding KG. Bacteria and wound healing. *Curr Opin Infect Dis* 2004;17:91–6.
- [29] Ovington L. Bacterial toxins and wound healing. *Ostomy Wound Manage* 2003;49:8–12.
- [30] Robson MC, Stenberg BD, Heggers JP. Wound healing alterations caused by infection. *Clin Plast Surg* 1990;17:485–92.
- [31] Laato M, Niinikoski J, Lundberg C, Gerdin B. Inflammatory reaction and blood flow in experimental wounds inoculated with *Staphylococcus aureus*. *Eur Surg Res* 1988;20:33–8.
- [32] Borkow G, Gabbay J. Copper as a biocidal tool. *Curr Med Chem* 2005;12:2163–75.
- [33] Hostynek JJ, Maibach HI. Copper hypersensitivity: dermatologic aspects – an overview. *Rev Environ Health* 2003;18:153–83.
- [34] Gorter RW, Butorac M, Cobian EP. Examination of the cutaneous absorption of copper after the use of copper-containing ointments. *Am J Ther* 2004;11:453–8.
- [35] Bilian X. Intrauterine devices. *Best Pract Res Clin Obstet Gynaecol* 2002;16:155–68.
- [36] Hubacher D, Lara-Ricalde R, Taylor DJ, Guerra-Infante F, Guzman-Rodriguez R. Use of copper intrauterine devices and the risk of tubal infertility among nulligravid women. *N Engl J Med* 2001;345:561–7.
- [37] Trumbo P, Yates AA, Schlicker S, Poos M. Dietary reference intakes: vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. *J Am Diet Assoc* 2001;101: 294–301.



NATURE + SCIENCE = BETTER WOUND CARE

MedCu's antimicrobial wound dressings harness the power of copper to set a new standard in wound care.

As a natural essential mineral for the human body, copper has been used for centuries to care for and treat wounds. Today, copper is at the forefront of a revolution in advanced wound care. MedCu is the first and only antimicrobial wound dressing impregnated with copper oxide microparticles to receive FDA clearance and CE Mark.

MedCu wound dressings offer protection against a broad spectrum of pathogens, including antibiotic-resistant bacteria. The dressings are non-adherent, with no need to pre-wet, and can be cut to ensure optimal fit for patient comfort. With sustained antimicrobial action, the dressings have up to 7 day wear-time allowing for fewer changes and reduced clinician contact time.

MedCu allows clinicians to effectively and efficiently care for wounds towards complete healing.

Areas of Use

MedCu's wound dressings are easy to apply and remove, and are suitable for a wide variety of applications including:

- Diabetic Wounds
- Leg & Foot Ulcers
- Pressure Ulcers
- First and Second-Degree Burns
- Surgical Wounds



SKU	Size inches	Size cm	Absorption Weight/Weight	Adhesive Contour
2C-0506-01	2x2.4	5x6	750%	-
2C-1012-01	4x4.5	10x12	1000%	-
2C-1020-01	4x8	10x20	1000%	-
2C-2020-01	8x8	20x20	1000%	-
2C-0505-01a	4x4 Pad: 2.5x2.5	10x10 Pad: 5x5	750%	+
2C-1025-01a	4x10 Pad: 4x7.8	10x25 Pad: 5x20	750%	+
3C-1012-01	4x4.5	10x12	800%	-